arXiv:quant-ph/0211133v1 20 Nov 2002

Imprinting a complete information about a quantum channel on its output state

Giacomo Mauro D'Ariano[*] and Paoloplacido Lo Presti[†]
Quantum Optics and Information Group, *Istituto Nazionale di Fisica della Materia, Unità di Pavia*[‡] *and Dipartimento di Fisica "A. Volta", via Bassi 6, I-27100 Pavia, Italy*
(Dated: February 1, 2008)

We introduce a novel property of bipartite quantum states, which we call *faithfulness*, and we say that a state is faithful when acting with a channel on one of the two quantum systems, the output state carries a complete information about the channel. The concept of faithfulness can also be extended to sets of states, when the output states patched together carry a complete imprinting of the channel.

PACS numbers: 03.65.Ta 03.67.-a

When a quantum system enters a quantum channel, its state transforms according to a linear, trace-preserving, and completely positive map $\mathcal{E}$ [1]. The input-output evolution of the state ρ can be written in a so-called Krauss's form [2]

$$\rho \rightarrow \mathcal{E}(\rho) = \sum_n K_n \rho K_n^\dagger \;, \tag{1}$$

where K_n are operators on the Hilbert space H of the quantum system, and satisfy the completeness relation $\sum_n K_n^\dagger K_n = I$, in order to preserve the trace of ρ. It is natural now to pose the question: is it possible to recover the channel $\mathcal{E}$ from the output state $\mathcal{E}(\rho)$?

It is clear that a single input state cannot be sufficient, and one actually needs to vary the input state over a somewhat "complete" set in order to recover the channel [3], and this is the basis of the so called *quantum process tomography* [4]. In Ref. [5] it has been shown how a fixed maximally entangled state can be used to recover the channel $\mathcal{E}$, with the entangled state playing the role of all possible input states in a *quantum parallel* fashion. The key feature of the method is that only one of the two entangled systems is impinged into the channel, whereas the other is left untouched, as in Fig. 1. The result is that for an unnormalized maximally entangled input, such as

$$|I\rangle\!\rangle = \sum_l |l\rangle \otimes |l\rangle, \tag{2}$$

where we used the notation $|A\rangle\!\rangle = \sum_{ij} A_{ij} |i\rangle \otimes |j\rangle$ for vectors $|A\rangle\!\rangle \in \mathsf{H} \otimes \mathsf{H}$ for a fixed orthonormal basis $\mathsf{b} = \{|i\rangle \otimes |j\rangle\}$ for $\mathsf{H} \otimes \mathsf{H}$, the corresponding unnormalized output

$$S_\mathcal{E} \doteq \mathcal{E} \otimes \mathcal{I}(|I\rangle\!\rangle\langle\!\langle I|) \tag{3}$$

is in one-to-one correspondence with the quantum channel $\mathcal{E}$, the inverse relation being

$$\mathcal{E}(\rho) = \mathrm{Tr}_2[I \otimes \rho^\tau \, S_\mathcal{E}], \tag{4}$$

where O^τ denotes transposition of the operator O with respect to the same basis b. The above one-to-one correspondence is an application of the Gelfand-Naimark-Segal representation (GNS) [6] of C*-algebras: a simple proof can be found in Ref. [7]. Notice that we have kept the input vector unnormalized, in order to have the correspondence valid also for infinite dimensions (the correspondence actually holds between completely bounded maps and bounded operators).



FIG. 1: The input state R is called *faithful* when the correspondence between the output state $R_\mathcal{E} \doteq \mathcal{E} \otimes \mathcal{I}(R)$ and the quantum channel $\mathcal{E}$ is one-to-one, namely the output state $R_\mathcal{E}$ carries a complete imprinting of the quantum channel $\mathcal{E}$.

The question is now: is it possible to keep the correspondence between output state $R_\mathcal{E}$ and channel $\mathcal{E}$ one-to-one also using a non maximally entangled input state R? As we will show in this letter, the answer is affirmative. Moreover, quite surprisingly there are even input *separable states* which work as well. We will call the input states with such property *faithful*, namely we will use the following definition: *a state R is* faithful *when the output $R_\mathcal{E} \doteq \mathcal{E} \otimes \mathcal{I}(R)$ from a quantum channel $\mathcal{E}$ is in one-to-one correspondence with $\mathcal{E}$.* In other words, $R_\mathcal{E}$ carries a complete imprinting of the quantum channel $\mathcal{E}$.

Using the spectral decomposition $R = \sum_l |A_l\rangle\!\rangle\langle\!\langle A_l|$ for the input state R, we can write

$$\begin{aligned} R_\mathcal{E} &= \mathcal{E} \otimes \mathcal{I}(R) = \sum_l I \otimes A_l^\tau S_\mathcal{E} I \otimes A_l^* \\ &\equiv \mathcal{E} \otimes \mathcal{R}(|I\rangle\!\rangle\langle\!\langle I|), \end{aligned} \tag{5}$$

where $\mathcal{R}(\rho) = \sum_l A_l^\tau \rho A_l^*$, and O^* denotes the complex conjugated of the operator O with respect to the basis b. From Eqs. (3) and (5) we can see that the *faithfulness of R is equivalent to the invertibility of the map $\mathcal{R}$*, since whenever the map $\mathcal{R}$ is invertible the output state $R_\mathcal{E}$ will be in one-to-one correspondence with $S_\mathcal{E}$, and thus with the channel $\mathcal{E}$. For a pure state $R \equiv |A\rangle\!\rangle\langle\!\langle A|$ faithfulness means simply that the operator A is invertible. In other

words: *a pure bipartite state is faithful iff it has maximal Schmidt's number*. More generally, for mixed states $R = \sum_l |A_l\rangle\rangle\langle\langle A_l|$, the full map $\mathcal{R}(\rho) = \sum_l A_l^\tau \rho A_l^*$ must be invertible, in order to have a one-to-one correspondence between $R_\mathcal{E}$ and $\mathcal{E}$.



FIG. 2: A generally mixed state $R = \sum_l |A_l\rangle\rangle\langle\langle A_l|$ is faithful when the map $\mathcal{R}(\rho) = \sum_l A_l^\tau \rho A_l^*$ is invertible, in order to guarantee the one-to-one correspondence between $R_\mathcal{E}$ and $\mathcal{E}$.

Let's first consider the case of finite dimensions, for simplicity. In terms of an orthonormal basis $\mathsf{B} = \{B_i\}$ of operators for the (Hilbert space of) operators $S \in \mathsf{B}(\mathsf{H})$ on H (i. e. $\mathrm{Tr}[B_i^\dagger B_j] = \delta_{ij}$), we have the unique expansion $S = \sum_i \mathrm{Tr}[B_i^\dagger S] B_i$. The map $\mathcal{R}$ is invertible iff it transforms the basis into a set of linearly independent operators $\mathcal{R}(B_i) = \sum_j R_{ij} B_j$, where, by definition, $R_{ij} = \mathrm{Tr}[B_j^\dagger \mathcal{R}(B_i)]$. In fact the map $\mathcal{R}$ is invertible iff the matrix R_{ij} has inverse R_{ij}^{-1}, and the inverse map $\mathcal{R}^{-1}$ is defined through its action on the basis as $\mathcal{R}^{-1}(B_i) = \sum_j R_{ij}^{-1} B_j$. Then, starting from $\mathcal{R}(S) = \sum_j \mathrm{Tr}[B_j^\dagger \mathcal{R}(S)] B_j$, it is possible to recover S by applying $\mathcal{R}^{-1}$ as $S = \sum_j \mathrm{Tr}[B_j^\dagger \mathcal{R}(S)] \mathcal{R}^{-1}(B_j)$. From the identity

$$R_{ij} = \mathrm{Tr}[B_j^\dagger \sum_l A_l^\tau B_i A_l^*] = \langle\langle B_j | \sum_l A_l^\tau \otimes A_l^\dagger | B_i \rangle\rangle, \quad (6)$$

we see that an equivalent condition for the invertibility of $\mathcal{R}$ is that the operator $\check{R}$ on $\mathsf{H}^{\otimes 2}$

$$\check{R} \doteq \sum_l A_l^\tau \otimes A_l^\dagger = (ER)^{\tau_2} E \equiv (R^{\tau_2} E)^{\tau_1} \quad (7)$$

is invertible, where $E = \sum_{ij} |ij\rangle\langle ji|$ is the swap operator, and O^{τ_l} denotes the partial transposition of the operator O on the lth Hilbert space. Actually, the correspondence $\mathcal{R} \leftrightarrow \check{R}$ preserves multiplication of maps, as $|\mathcal{R}(S)\rangle\rangle = \check{R}|S\rangle\rangle$, whence $|\mathcal{R}^{-1}(S)\rangle\rangle = \check{R}^{-1}|S\rangle\rangle$. This further clarifies the relation between the invertibility of the map $\mathcal{R}$ and that of the operator $\check{R}$ [8].

Therefore, a bipartite state R is *faithful* if and only if the operator $\check{R}$ in Eq. (7) is invertible. For this kind of states the relation $R_\mathcal{E} = \mathcal{E} \otimes \mathcal{I}(R) \leftrightarrow S_\mathcal{E}$ is one-to-one, and the information about the channel $\mathcal{E}$ can be extracted from $R_\mathcal{E}$ as follows

$$\mathcal{E}(\rho) = \mathrm{Tr}_2[I \otimes \rho^\tau \mathcal{I} \otimes \mathcal{R}^{-1}(R_\mathcal{E})]. \quad (8)$$

It is clear that the set of faithful states R is *dense* within the set of all bipartite states. However, the knowledge of the map $\mathcal{E}$ from a measured $R_\mathcal{E}$ will be affected by increasingly large errors for $\check{R}$ approaching a non-invertible operator, and *measures of faithfulness* should be introduced (we will analyze this problem later in this letter). Since the set of faithful states is dense, it follows that there must be faithful states among mixed separable states. For example, the Werner's states for dimension d

$$R_f = \frac{1}{d(d^2-1)}[(d-f) + (df-1)E], \quad -1 \le f \le 1, \quad (9)$$

are separable for $f \ge 0$, however, they are faithful for all $f \neq \frac{1}{d}$, since one has $(ER_f)^{\tau_2} = \frac{1}{d(d^2-1)}[(d-f)|I\rangle\rangle\langle\langle I| + (df-1)]$, and the singular values of $\check{R}_f$ are $\frac{df-1}{d(d^2-1)}$ and $\frac{1}{d}$. Similarly, the "isotropic" states

$$R_f = \tfrac{f}{d}|I\rangle\rangle\langle\langle I| + \tfrac{1-f}{d^2-1}(I - \tfrac{1}{d}|I\rangle\rangle\langle\langle I|), \quad (10)$$

are faithful for $f \neq \frac{1}{d^2}$ and separable for $f \le \frac{1}{d}$, the singular values of $\check{R}_f$ being $\frac{d^2 f-1}{d(d^2-1)}$ and $\frac{f}{d}$.

For infinite dimensions (e. g. for "continuous variables"), one needs to restrict $\mathsf{B}(\mathsf{H})$ to the Hilbert space of Hilbert-Schmidt operators on H, and this lead to the problem that the inverse map $\mathcal{R}^{-1}$ is unbounded. The result is that we will recover the channel $\mathcal{E}$ from the measured $R_\mathcal{E}$, however, with unbounded amplification of statistical errors, depending on the chosen representation $\mathsf{B} = \{B_j\}$, due to the fact that $\mathcal{R}^{-1}(B_j)$ increases unboundedly for $j \to \infty$. As an example, let's consider a twin beam from parametric down-conversion of vacuum

$$|\Psi\rangle\rangle = \Psi \otimes I|I\rangle\rangle, \quad \Psi = (1-|\xi|^2)^{\frac{1}{2}} \xi^{a^\dagger a}, \qquad |\xi| < 1 \quad (11)$$

for a fixed ξ, $a^\dagger$ and a, with $[a, a^\dagger] = 1$, denoting the creation and annihilation operators of the harmonic oscillator describing the field mode corresponding to the first Hilbert space in the tensor product (in the following we will denote by $b^\dagger$ and b the creation and annihilation operators of the other field mode). The state is faithful, but the operator Ψ^{-1} is unbounded, whence the inverse map $\mathcal{R}^{-1}$ is also unbounded. In a photon number representation $\mathsf{B} = \{|n\rangle\langle m|\}$, the effect will be an amplification of errors for increasing numbers n, m of photons.

Consider now the quantum channel describing the *Gaussian displacement noise* [13]

$$\mathcal{N}_\nu(\rho) = \int_\mathbb{C} \frac{\mathrm{d}\,\alpha}{\pi\nu} \exp[-|\alpha|^2/\nu] D(\alpha)\rho D^\dagger(\alpha), \quad (12)$$

where $D(\alpha) = \exp(\alpha a^\dagger - \alpha^* a)$ denotes the usual displacement operator on the phase space. The Gaussian noise is in a sense the analogous of the depolarizing channel for infinite dimension. The maps $\mathcal{N}_\nu$ for varying ν satisfy the multiplication rule $\mathcal{N}_\nu \mathcal{N}_\mu = \mathcal{N}_{\nu+\mu}$, whence the inverse map is formally given by $\mathcal{N}_\nu^{-1} \equiv \mathcal{N}_{-\nu}$. Notice that, since the map $\mathcal{N}_\nu$ is compact, the inverse map $\mathcal{N}_\nu^{-1}$ is

necessarily unbounded. As a faithful state consider now the mixed state given by the twin-beam, with one beam spoiled by the Gaussian noise, namely

$$R = \mathcal{I} \otimes \mathcal{N}_\nu(|\Psi\rangle\!\rangle\langle\!\langle\Psi|). \tag{13}$$

A lengthy straightforward calculation gives the state

$$R = \frac{1}{\nu}\Psi \otimes I \exp[-(a - b^\dagger)(a^\dagger - b)/\nu]\Psi^\dagger \otimes I, \tag{14}$$

and its partial transposed

$$R^{\tau_2} = (\nu+1)^{-1}\Psi \otimes I \left(\frac{\nu-1}{\nu+1}\right)^{\frac{1}{2}(a-b)^\dagger(a-b)} \Psi^\dagger \otimes I, \tag{15}$$

where transposition is defined with respect to the basis of eigenvectors of $a^\dagger a$ and $b^\dagger b$. Since our state is Gaussian, the PPT criterion guarantees separability [14], and for $\nu > 1$ our state (14) is separable (see also Ref. [15]), still it is *formally* faithful, since the operator Ψ and the map N_ν are both invertible. Notice that unboundedness of the map inversion can even wash out completely the information on the channel in some particular chosen representation $\mathsf{B} = \{B_j\}$, e. g. when all operators B_j are out of the boundedness domain of $\mathcal{R}^{-1}$. This is the case, for example, of the (overcomplete) representation $\mathsf{B} = \{|\alpha\rangle\langle\beta|\}$, with $|\alpha\rangle$ and $|\beta\rangle$ coherent states, since from the identity

$$\mathcal{N}_\nu(|\alpha\rangle\langle\alpha|) = \frac{1}{\nu+1}D(\alpha)\left(\frac{\nu}{\nu+1}\right)^{a^\dagger a} D^\dagger(\alpha), \tag{16}$$

one obtains

$$\mathcal{N}_\nu^{-1}(|\alpha\rangle\langle\alpha|) = \frac{1}{1-\nu}D(\alpha)\left(1-\nu^{-1}\right)^{-a^\dagger a} D^\dagger(\alpha), \tag{17}$$

which has convergence radius $\nu \le \frac{1}{2}$, which is the well known bound for Gaussian noise for the quantum tomographic reconstruction for coherent-state and Fock representations [16]. Therefore, we say that the state is *formally* faithful, however, we are constrained to representations which are analytical for the inverse map $\mathcal{R}^{-1}$.

In a more general framework, we can consider the faithfulness of the bipartite state R of two quantum systems described by different (in finite dimensions) Hilbert spaces H and K. We need now to consider operators A in either $\mathsf{B}(\mathsf{K},\mathsf{H})$, $\mathsf{B}(\mathsf{H})$, or $\mathsf{B}(\mathsf{K})$ (in all cases we will keep the same notation $|A\rangle\!\rangle$ for the corresponding vector in $\mathsf{H}\otimes\mathsf{K}$, $\mathsf{H}^{\otimes 2}$, or $\mathsf{K}^{\otimes 2}$). Corresponding to the state $R = \sum_l |A_l\rangle\!\rangle\langle\!\langle A_l|$ on $\mathsf{H}\otimes\mathsf{K}$ we have now the map $\mathcal{R}(\rho) = \sum_l A_l^\tau \rho A_l^*$ from states on H to states on K, and the related operator $\check{R} = \sum_l A_l^\tau \otimes A_l^\dagger \in \mathsf{B}(\mathsf{H}^{\otimes 2},\mathsf{K}^{\otimes 2})$. We still have $\check{R}|S\rangle\!\rangle = |\mathcal{R}(S)\rangle\!\rangle$, with $|S\rangle\!\rangle \in \mathsf{H}^{\otimes 2}$, and $|\mathcal{R}(S)\rangle\!\rangle \in \mathsf{K}^{\otimes 2}$. In order to express $\check{R}$ through its matrix components R_{ij}, we now need to choose d an operator basis $\mathsf{B}' = \{B'_i\}$ also for $\mathsf{B}(\mathsf{K})$, so that $\mathcal{R}(B_i) = \sum_j R_{ij}B'_j$, where now

$$R_{ij} = \mathrm{Tr}[B_j'^\dagger \mathcal{R}(B_i)] = \langle\!\langle B'_j|\sum_l A_l^\tau \otimes A_l^\dagger|B_i\rangle\!\rangle\ . \tag{18}$$

Then, the faithfulness of R, is more generally equivalent to the left invertibility of $\mathcal{R}$, or, equivalently, to the left invertibility of $\check{R}$.

Now, if we have a set of bipartite states $\{R^{(n)}\}_1^N$ on $\mathsf{H}\otimes\mathsf{K}$ which are not faithful, we can try anyway to recover the quantum channel $\mathcal{E}$ by patching the outputs $R_{\mathcal{E}}^{(n)} \doteq \mathcal{E}\otimes\mathcal{I}(R^{(n)})$ together. This is possible if and only if the following joint state on $\mathsf{H}\otimes\tilde{\mathsf{K}}$ ($\tilde{\mathsf{K}} \doteq \mathsf{K}\otimes\mathbb{C}^N$) is faithful

$$R_{\rm set} = \sum_{n=1}^N p_n R^{(n)} \otimes |n\rangle\langle n|, \tag{19}$$

where p_n are probabilities (the state $R_{\rm set}$ is equivalent to a mixture the states $\{R^{(n)}\}$ in a *knowingly* fashion). In this case we call the set $\{R^{(n)}\}$ a *faithful set of states*.

What can we do with an unfaithful state R? The map $\mathcal{R}$ is not invertible, and for all vectors $|S\rangle\!\rangle \in \mathsf{Ker}(\check{R})$ we have $\mathcal{R}(S) = 0$. However, an unfaithful state can still be useful in recovering some quantum channels, or at least a part of them. In fact, one can use the pseudo-inverse $\check{R}^\ddagger$, which allows inversion only in $\mathsf{Rng}(\check{R}^\dagger) \equiv \mathsf{Ker}(\check{R})^\perp$, with $\check{R}^\ddagger\check{R} = I - P$, P being the orthogonal projector on $\mathsf{Ker}(\check{R})$. Correspondingly, one defines the pseudo-inverse map $\mathcal{R}^\ddagger$ from the identity $|\mathcal{R}^\ddagger(S)\rangle\!\rangle \doteq \check{R}^\ddagger|S\rangle\!\rangle$, or, equivalently, by its action on the basis for $\mathsf{B}(\mathsf{K})$, i. e. $\mathcal{R}^\ddagger(B'_i) = \sum_j R^\ddagger_{ij}B_j$. It is clear that now the inversion, instead of the full $S_{\mathcal{E}}$, will give its projection

$$\tilde{S}_{\mathcal{E}} = \mathcal{I}\otimes\mathcal{R}^\ddagger(R_{\mathcal{E}}) = \mathcal{I}\otimes\mathcal{Q}(S_{\mathcal{E}}) \tag{20}$$

where $\mathcal{Q} = \mathcal{R}^\ddagger\mathcal{R} = \mathcal{Q}^2$ is the orthogonal projection map on the operator space $\mathsf{B}(\mathsf{H})$, also defined as $|\mathcal{Q}(S)\rangle\!\rangle = (I-P)|S\rangle\!\rangle = \check{Q}|S\rangle\!\rangle$. The partially recovered map $\tilde{\mathcal{E}}(\rho) = \mathrm{Tr}_2[I\otimes\rho^\tau\tilde{S}_{\mathcal{E}}]$ is generally not CP, and can also be written as $\tilde{\mathcal{E}} = \mathcal{E}\mathcal{Q}^*$, where $\mathcal{Q}^*$ is the orthogonal projection map corresponding to the operator $\check{Q}^*$.

The above considerations suggest a definition of a *number of faithfulness* φ as $\varphi(R) = \mathrm{Tr}(\check{Q})$, corresponding to the rank of $\check{R}$. Then, a state is faithful for $\varphi(R) = \dim(\mathsf{H})^2$. Notice that for $\varphi(R) < \dim(\mathsf{H})^2$ one can have the situation in which $\mathsf{Rng}(\check{R}^\dagger) = \mathsf{Span}\{|S\rangle\!\rangle, S\,\text{commuting}\}$, in which case the state R allows to reconstruct completely only “classical” channels, with the input restricted to an abelian algebra of states.

The introduction of pseudo-inversion provides an algorithm for the patching when using a set of states $\{R^{(n)}\}$ that lead to the projection maps $\{\mathcal{Q}^{(n)}\}$. The set is faithful iff we can recover any operator $S \in \mathsf{B}(\mathsf{H})$ from its projections $\mathcal{Q}^{(n)}(S)$, and this is possible iff, given a basis $\{B_i\}$ for $\mathsf{B}(\mathsf{H})$, one has $\mathsf{Span}\{\mathcal{Q}^{(n)}(B_i)\}_{i,n} = \mathsf{B}(\mathsf{H})$. In such circumstances, any element of the basis can be expressed as a linear combination of the $\mathcal{Q}^{(n)}(B_i)$, therefore the component $\mathrm{Tr}[B_i^\dagger S]$ of S will be a linear combination of terms of the form $\mathrm{Tr}[\mathcal{Q}^{(n)}(B_i)^\dagger S]$, whereas these terms can be calculated substituting S with the projection $\mathcal{Q}^{(n)}(S)$. Expanding $B_i = \sum_{jn}\lambda^n_{ij}\mathcal{Q}^{(n)}(B_j)$,

the reconstruction formula for $S_{\mathcal{E}}$ will be

$$S_{\mathcal{E}} = \sum_{ijn} \lambda_{ij}^{n*} \operatorname{Tr}_2[I \otimes \mathcal{Q}^{(n)}(B_j)^\dagger \tilde{S}_{\mathcal{E}}^{(n)}] \otimes B_i \ . \tag{21}$$

Now, let's consider the problem of how to define a measure of faithfulness $F(R)$ of the state R. As already noticed, even though in principle any faithful state can be used to perform a tomography of the channel $\mathcal{E}$ [5], the experimental errors on the measured $R_{\mathcal{E}}$ are propagated to $\mathcal{E}$ by the inversion of the map $\mathcal{R}$. Thus different faithful input states can produce very different errors on the measured channel. It is clear that all the features producing amplification of errors are contained in the singular values σ_l of $\check{R}$: the inversion involves multiplications by σ_l^{-1}, then the smaller are the singular values σ_l, the larger is the error amplification, whence the number of measurements needed to have a good reconstruction. This suggests that a measure of faithfulness should be an increasing function $F(R) \equiv f(\{\sigma_l\})$ of the singular values σ_l of $\check{R}$, and when the different σ_l are equivalent $f(\{\sigma_l\})$ must be invariant under their permutations, whence $F(R)$ is a symmetric monotone norm of $\check{R}$ [17]. It is clear, however, that it is unpractical to have a universal measure for faithfulness, and its actual definition will be dictated by the goodness criterion adopted for the reconstruction of the quantum channel $\mathcal{E}$. For example, the Frobenius norm $\|\check{R}\|_2 = \operatorname{Tr}[\check{R}^\dagger \check{R}] = \operatorname{Tr}[R^\dagger R]$ coincides with the purity of the state R, and this simply shows that, for the part of the channel that can be reconstructed, the error is minimized for pure input state R.

The definitions of F and φ can be naturally extended to sets of states $\{R^{(n)}\}$ via the introduction of the joint state R_{set} in Eq. (19). It obviously follows that any chosen degree of faithfulness $F(R)$ of a maximally entangled pure state R will be larger than the degree $F(\{R^{(n)}\})$ of any faithful set $\{R^{(n)}\}$ of unfaithful states.

In conclusion, in this letter we have introduced a new feature of bipartite quantum states, which we call *faithfulness*, corresponding to the ability of the state of carrying the complete imprinting of a channel acting on one of the pair of quantum systems. This property has also been extended to sets of bipartite states, when the channel can be recovered from the corresponding output states patched together. We have seen that there are separable states that are faithful, and the maximally faithful states are the maximally entangled pure states. We want to stress that the property of being faithful is a strictly quantum feature, since a faithful state cannot be written as the mixture of local classical (i. e. commuting) states. This also shows how subtle is the game between the classical and quantum natures in the correlations of a general mixed quantum state.

One of us (G. M. D.) acknowledges B. Kűmmerer for pointing him the GNS and Kolmogorov's constructions, and P. Horodecki for interesting discussions. This work has been jointly founded by the EC under the program ATESIT (Contract No. IST-2000-29681), by the US Department of Defense MURI program (Grant No. DAAD19-00-1-0177), and by INFM project PRA-2002-CLON.

* Electronic address: dariano@unipv.it; Also at: Department of Electrical and Computer Engineering, Northwestern University, Evanston, IL 60208

† Electronic address: lopresti@unipv.it

‡ URL: `http://www.qubit.it`

[1] A map $\mathcal{E}$ is completely positive if it preserves positivity when acting locally on a bipartite state. In other words, upon denoting by $\mathcal{I}$ the identical map on the Hilbert space K of a second quantum system, the extended map $\mathcal{E} \otimes \mathcal{I}$ on $\mathsf{H} \otimes \mathsf{K}$ must be positive for any extension K.

[2] K. Kraus, *States, Effects, and Operations* (Springer-Verlag, Berlin, 1983)

[3] G. M. D'Ariano and L. Maccone, Phys. Rev. Lett. **80** 5465 (1998)

[4] I. L. Chuang and M. A. Nielsen, J. Mod. Opt. **44** 2455 (1997)

[5] G. M. D'Ariano, and P. Lo Presti, Phys. Rev. Lett. **86** 4195 (2001)

[6] G. J. Murphy, *C^*-Algebras and Operator Theory* (Academic Press, London, San Diego 1990)

[7] G. M. D'Ariano, and P. Lo Presti, Phys. Rev. A **64** 042308 (2001)

[8] Differently from the correspondence (3), the isomorphism $\check{R} \leftrightarrow \mathcal{R}$ doesn't preserve positivity, but preserves the algebraic structure of bounded maps[9]. Other isomorphisms can be obtained using the discrete 24 elements group [10] generated by the involutions τ_1, τ_2, right and left multiplications by E, adjoint, and partial adjoint. A relevant example is given by the *matrix realignement* of Refs. [11] and [12].

[9] G. M. D'Ariano, unpublished

[10] C. J. Oxenrider and R. D. Hill, Lin. Alg. Appl. **69** 205 (1985)

[11] O. Rudolph, quant-ph/0202121 (2002)

[12] M. Horodecki, P. Horodecki, and R. Horodecki, LANL arXive eprints quant-ph/0206008 (2002)

[13] M. J. W. Hall, Phys. Rev. A **50**, 3295 (1994)

[14] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000)

[15] M. G. A. Paris, J. Opt. B: Quantum Semiclass. Opt. **4** (2002) 442

[16] G. M. D'Ariano, and N. Sterpi, J. Mod. Optics **44** 2227 (1997)

[17] R. Bhatia, *Matrix Analysis* (Springer-Verlag, New York (1997)